

SHORT CONTRIBUTION TO THE GENUS *PACHYRHYNCHUS* FROM THE PHILIPPINES (COLEOPTERA: CURCULIONIDAE: PACHYRHYNCHINI)

Jiří Háva, Anita Rukmane

Háva J., Rukmane A. Short contribution to the genus *Pachyrhynchus* from the Philippines (Coleoptera: Curculionidae: Pachyrhynchini). *Acta Biol. Univ. Daugavp.*, 18 (2): 203 – 205.

The following species are newly recorded from the Philippines: *Pachyrhynchus apicatus* Schultze, 1922 (Luzon); *Pachyrhynchus occidentalis* Rukmane, 2017 (Visayas); *Pachyrhynchus rukmanae* Barševskis, 2016 (Luzon); *Pachyrhynchus speciosus* Waterhouse, 1841 (Visayas).

Key words: faunistics, new records, Coleoptera, Curculionidae, Pachyrhynchus, the Philippines.

Jiří Háva. Private Entomological Laboratory and Collection, Rýznerova 37, CZ - 252 62 Únětice u Prahy, Praha-západ, Czech Republic; e-mail: jh.dermeestidae@volny.cz.

Department of Forest Protection and Entomology, Faculty of Forestry and Wood Sciences, Czech University of Life Sciences, Kamýcká 129, CZ-165 00, Prague 6 - Suchbát, Czech Republic.

Anita Rukmane. Daugavpils University, Institute of Life Sciences and Technology, Coleopterological Research Center, Vienības Str. 13, Daugavpils, LV - 5401, Latvia.

INTRODUCTION

The genus *Pachyrhynchus* Germar, 1824 belong to subfamily Entiminae recently contains 145 species and subspecies (Rukmane 2018). The genus is well known from Oriental Region, with predomination in The Philippines (93%) and few species distributed in Indonesia (4%) and Japan (3%) (Rukmane 2018). In the present short article four species are newly recorded.

MATERIAL AND METHODS

The mentioned specimens deposited in collections:

JHAC - Private Entomological Laboratory & Collection, Únětice u Prahy, Prague-west, Czech Republic;

DUBC - Beetles Collection of Coleopterological Research Center, Institute of Life Sciences and Technology, Daugavpils University (Daugavpils, Latvia).

All material determined by second author Anita Rukmane.

RESULTS

Pachyrhynchus apicatus Schultze, 1922

Material examined: Philippines, West Luzon, Aurora, Dingalan, Sierra Madre, June 2018, local coll., 2 spec., (JHAC); June 2014, 1 spec., July 2014, 1 spec., September 2014, 1 spec, November 2014, 1 spec., June 2015, 1 spec., November 2015, 1 spec., April 2016, 1 spec., May 2016, 1. spec, all by local collectors (DUBC).

Total: 10 specimens.

Distribution. Species known from Philippines: Polillo, new for Luzon.

***Pachyrhynchus occidentalis* Rukmane, 2017**

Material examined: Philippines, Visayas, Samar, Lope de Vega, June 2018, local coll., 1 spec., (JHAC).

Distribution. Species known from Philippines: Mindanao, new for Visayas.

***Pachyrhynchus rukmanee* Barševskis, 2016**
(Fig. 1)

Material examined: Philippines, Luzon, II. 2017, local coll., 1 spec., (JHAC).

Distribution. Species known from Philippines: Marinduque, new for Luzon.

***Pachyrhynchus speciosus* Waterhouse, 1841**

Material examined: Philippines, Visayas, Samar, Hinabangan, March 2018, local coll., 1 spec., (JHAC); May 2013, 1 spec., August 2013, 1 spec., October 2015, 2 spec., January 2016, 1 spec., May 2016, 1 spec., June 2016, 1 spec., October 2017, 2 spec., all by local collectors (DUBC).

Total: 10 specimens.

Distribution. Species known from Philippines: Mindanao, new for Visayas.

DISCUSSION

Current study showed new distribution range for four species of genus *Pachyrhynchus*.

Pachyrhynchus apicatus Schultze, 1922 which were previously recorded only from Pollilo island occur also on Luzon Island. Another case of such distribution range were previously recorded only for *Pachyrhynchus moniliferus* Germar, 1824 (Schultze 1923). Repetitive cases of various species abundant on both islands reveals new

possible distribution pathway for species of genus *Pachyrhynchus*, further investigation of species common for both Pollilo and Luzon Islands is required.

Pachyrhynchus occidentalis Rukmane, 2017 which were previously recorded only from Mindanao Island occur also on Samar Island. In various articles about species distribution pathways authors refer to greater Mindanao or Mindanao PAIC (pleistocene aggregate island complex) species distribution. Similarly with *Pachyrhynchus speciosus* Waterhouse, 1841 which according to new distribution data is also distributed on Mindanao and Samar Islands.



Fig.1. Dorsal habitus of: A - *P. rukmanee*.

Pachyrhynchus rukmanee Barševskis, 2016 which were previously recorded only from Marinduque Island occur also on Luzon Island. This species exhibits allopatric convergence in color and markings with *Pachyrhynchus masatoshii* Yoshitake & Yap, 2018 from Luzon Island.

Accepted: 01.12.2018.

On account of flightless abilities and attraction to mountain ecosystems, weevils of genus *Pachyrhynchus* show high level of endemism and one particular species tend to inhabit single island, which is the case for majority of species in genus. Nevertheless, previous studies revealed that some species tend to occur on various islands, for example *P. erichsoni* Waterhouse, 1841 which is abundant on both Luzon and Mindanao (Schultze 1923). Current study is one more assertion.

REFERENCES

- Rukmane A. 2017. New species of the genus *Pachyrhynchus* Germar (Coleoptera: Curculionidae: Entiminae) from the Greater Mindanao Pleistocene Aggregate Island complex (Philippines). *Acta Biologica Universitates Daugavpiliensis*, 17(1): 85-95
- Rukmane A. 2018. Annotated checklist of the genus *Pachyrhynchus* (Coleoptera: Curculionidae: Pachyrhynchini). *Acta Biologicae Universitates Daugavpiliensis*, 18(1): 63-68.
- Schultze W. 1923. A monograph of the pachyrhynchid group of the Brachyderinae, Curculionidae: part 1. *The Philippine Journal of Science*, (23): 609-673.

Received: 03.11.2018.